

To appear in the September 21, 2000 issue of Nature

The Accelerations of Stars Orbiting the Milky Way's Central Black Hole

A. M. Ghez*, M. Morris*, E. E. Becklin*, A. Tanner*, & T. Kremenek*

*Dept. of Physics and Astronomy, UCLA, Los Angeles, CA 90095-1562

Recent measurements¹⁻⁴, of the velocities of stars near the center of the Milky Way have provided the strongest evidence for the presence of a supermassive black hole in a galaxy⁵, but the observational uncertainties poorly constrain many of the properties of the black hole. Determining the accelerations of stars in their orbits around the center provides much more precise information about the position and mass of the black hole. Here we report measurements of the accelerations for three stars located ~ 0.005 pc (projected on the sky) from the central radio source Sagittarius A* (Sgr A*); these accelerations are comparable to those experienced by the Earth as it orbits the Sun. These data increase the inferred minimum mass density in the central region of the Galaxy by an order of magnitude relative to previous results and localized the dark mass to within 0.05 ± 0.04 arcsec of the nominal position of Sgr A*. In addition, the orbital period of one of the observed stars could be as short as 15 years, allowing us the opportunity in the near future to observe an entire period.

In 1995, we initiated a program of high resolution $2.2 \mu m$ (K band) imaging of the inner $5 \text{ arcsec} \times 5 \text{ arcsec}$ of the Galaxy's central stellar cluster with the W. M. Keck 10-m telescope on Mauna Kea, Hawaii ($1 \text{ arcsec} = 0.04 \text{ pc}$ at the distance to the Galactic Center, 8 kpc, ref. 6). From each observation, several thousand short exposure ($t_{exp}=0.137$

sec) frames were collected, using the facility near-infrared camera, NIRC^{7,8}, and combined to produce a final diffraction-limited image, having an angular resolution of 0.05 arcsec. Between 1995 and 1997, images were obtained once a year with the aim of detecting the stars’ velocities in the plane of the sky. The results from these measurements are detailed in ref. 4; in summary, two-dimensional velocities were measured for 90 stars with simple linear fits to the positions as a function of time. These velocities, which reach up to 1400 km/sec, implied the existence of a $2.6 \times 10^6 M_{\odot}$ black hole coincident (± 0.1 arcsec) with the nominal location of Sgr A* (ref. 9), the unusual radio source^{10,11,12} long believed to be the counterpart of the putative black hole.

The new observations presented here were obtained several times a year from 1997 to 1999 to improve the sensitivity to accelerations. With nine independent measurements, we now fit the positions of stars as a function of time with second order polynomials (see Fig. 1). The resulting velocity uncertainties are reduced by a factor of 3 compared to our earlier work, primarily as a result of the increased time baseline, and, in the central square arcsecond, by a factor of 6 that presented in ref. 3, due to our higher angular resolution. Among the 90 stars in our original proper motion sample⁴, we have now detected significant accelerations for three stars, S0-1, S0-2 and S0-4 (see Table 1); specifically, these are the sources for which the reduced chi-squared for a quadratic fit is smaller than that for the linear model of their motions by more than 1. These three stars are independently distinguished in our sample, being among the fastest moving stars ($v = 560$ to 1350 km/sec) and among the closest to the nominal position of Sgr A* ($r_{1995} = 0.004$ to 0.013 pc). With accelerations of 2-5 milli-arcsecond/yr², or equivalently $(3 - 6) \times 10^{-6}$ km/sec², they are experiencing accelerations similar to the Earth in its orbit about the Sun.

Acceleration vectors, in principle, are more precise tools than the velocity vectors for studying the central mass distribution. Even projected onto the plane of the sky, each

acceleration vector should be oriented in the direction of the central mass, assuming a spherically symmetric potential. Thus, the intersection of multiple acceleration vectors is the location of the dark mass. Figure 2 shows the acceleration vector’s direction for the three stars. Within 1σ , these vectors do indeed overlap, and furthermore, the intersection point lies a mere 0.05 ± 0.03 arcsec East and 0.02 ± 0.03 arcsec South of the nominal position of Sgr A*, consistent with the identification of Sgr A* as the carrier of the mass. Previously, with statistical treatments of velocities only, the dynamical center was located to within ± 0.1 arcsec (1σ) of Sgr A*’s position⁴. This velocity-based measurement is unaffected by the increased time baseline, as its uncertainty is dominated by the limited number of stars at a given radius. Therefore, the accelerations improve the localization of our Galaxy’s dynamical center by a factor of 3, which is critical for reliably associating any near-infrared source with the black hole given the complexity of the region.

Like the directions, the magnitudes of the acceleration vectors also constrain the central black hole’s properties. In three dimensions, the acceleration and radius vectors (a_{3-D} and r_{3-D} , respectively) provide a direct measure of the enclosed mass simply by $a_{3-D} = G \times M/r_{3-D}^2$. For a central potential, the acceleration and radius vectors are co-aligned and, with a projection angle to the plane of the sky θ , the two dimensional projections place the following lower limit on the central mass: $M \cos^3(\theta) = a_{2-D} \times r_{2-D}^2 / G$. This analysis is independent of the star’s orbital parameters, although θ is in fact a lower limit for the orbital inclination angle. Figure 3 shows the minimum mass implied by each star’s two-dimensional acceleration as a function of projected radius, along with dashed curves displaying how the implied mass and radius grow with projection angle. For each point, the uncertainty in the position of the dynamical center dominates the minimum mass uncertainties. Also plotted are the results from the statistical analysis of the velocity vectors measured in the plane of the sky, which imply a dark mass of $2.3 - 3.3 \times 10^6 M_\odot$ inside a radius of 0.015 pc^{3,4}. If, as has been assumed, this dark mass is in the form of

a single supermassive black hole, the enclosed mass should remain level at smaller radii. Projection angles of 51 - 56 and 25 - 37 degrees for S0-1 and S0-2, respectively, would yield the mass inferred from velocities and place these stars at a mere ~ 0.008 pc (solid blue portions of the limiting mass curves in Figure 3), thus increasing the dark mass density implied by velocities by an order of magnitude to $8 \times 10^{12} M_{\odot}/pc^3$. With smaller projection angles, these two stars also allow for the enclosed mass to decrease at smaller radii, as would occur in the presence of an extended distribution of dark matter surrounding a less massive black hole^{13–17}. In contrast to the agreement between the mass distribution inferred from the velocities and the accelerations for S0-1 and S0-2, the minimum mass implied by S0-4's acceleration is inconsistent at least at the 1σ level. We note, in support of the validity of S0-4's acceleration vector, that its orientation is consistent with the intersection of the S0-1 and S0-2 acceleration vectors (see Fig. 2). Nonetheless, continued monitoring of S0-4 will be important for assessing this possible discrepancy. Overall, the individual magnitudes of the three acceleration vectors support the existence of a central black hole with mass $\sim 3 \times 10^6 M_{\odot}$.

With acceleration measurements, it is now possible to constrain the individual orbits. We checked the conclusions of the previous studies^{3,4} by assuming that the stars are bound to a central mass of $2.3 - 3.3 \times 10^6 M_{\odot}$ located within 0.03 arcsec of the nominal position of Sgr A*. Excellent orbital fits were found for S0-1 and S0-2 for the entire range of masses and for true focii within 0.01 arcsec of the nominal position of Sgr A*, suggesting that we now have comparable accuracy in determining the IR location of Sgr A* (ref. 9) and the dynamical center of our Galaxy (see Fig. 4). The orbital solutions for these stars have eccentricities ranging from 0 (circular) to 0.9 for S0-1 and 0.5 to 0.9 for S0-2 and periods ranging from 35 - 1,200 and 15 - 550 years, respectively, raising the possibility of seeing a star make a complete journey around the center of the Galaxy within the foreseeable future. Although the fits are not yet unique, they impose a maximum orbital distance

from the black hole, or apoapse, of 0.1 pc. This suggests that S0-1 and S0-2 might have formed locally. If these stars are indeed young², then their small apoapse distance presents a challenge to classical star formation theories in light of the strong tidal forces created by the central black hole and might require a collisional or compressional star formation scenario^{18,19}. However, dynamical friction may be able to act on a short enough time scale to bring these stars in from a much larger distance²⁰. More accurate orbital parameters are needed to fully address these problems and others such as the distance to the Galactic Center²¹. The determinations of these parameters will be considerably improved when radial velocities are measured for these stars using adaptive optics techniques, as the current solutions based on the proper motion data alone predict radial velocities ranging from 200 to 2000 km/sec.

REFERENCES

1. Eckart, A. & Genzel, R. Stellar proper motions in the central 0.1 pc of the Galaxy. *Mon. Not. R. Astron. Soc.*, **284**, 576-598 (1997)
2. Genzel, R., Eckart, A., Ott, T. and Eisenhauer, F. On the nature of the dark mass in the centre of the Milky Way. *Mon. Not. R. Astron. Soc.*, **291-234**, 219 (1997)
3. Genzel, R., Pichon, C., Eckart, A., Gerhard, O.E., and Ott, T. Stellar Dynamics in the Galactic centre: Proper Motions and Anisotropy. *Mon. Not. R. Astron. Soc.*, in press (2000)
4. Ghez, A. M., Klein, B. L., Morris, M., and Becklin, E. E. High Proper-Motion Stars in the Vicinity of Sagittarius A*: Evidence for a Supermassive Black Hole at the Center of Our Galaxy, *Astrophys. J.*, **509**, 678-686 (1998)
5. Maoz, E. 1998, Dynamical Constraints on Alternatives to Supermassive Black Holes in Galactic Nuclei. *Astrophys. J.*, **494**, L181-L184 (1998)
6. Reid, M. J. The distance to the center of the Galaxy. *Ann. Rev. Astron. Astrophys.* **31**, 345-372 (1993)
7. Matthews, K. and Soifer, B. T. The Near Infrared Camera on the W. M. Keck Telescope. in *Astronomy with Infrared Arrays: The Next Generation* (ed. I. McLean) 239-246 (Kluwer Academic Publications, Dordrecht, Astrophysics and Space Sciences Library, **190**, 1994)
8. Matthews, K., Ghez, A. M., Weinberger, A. J., and Neugebauer, G. The First Diffraction-Limited Images from the W. M. Keck Telescope. *Proc. Astron. Soc. Pac.*, **108**, 615-619 (1996)
9. Menten, K. M., Reid, M. J., Eckart, A., & Genzel, R. The Position of Sgr A*: Accurate Alignment of the Radio and Infrared Reference Frames at the Galactic Center.

- Astrophys. J.*, **475**, L111-L114 (1997)
10. Lo, K.Y., Shen, Z.-Q., Zhao, J.-H., & Ho, P.T. Intrinsic Size of SGR A*: 72 Schwarzschild Radii. *Astrophys. J.*, **508**, L61-L64 (1998)
 11. Reid, M.J., Readhead, A.C.S, Vermeulen, R.C., Treuhaft, R.N. The Proper Motion of Sagittarius A*. I. First VLBA Results. *Astrophys. J.*, **524**, 816-823 (1999)
 12. Backer, D.C., & Sramek, R.A. Proper Motion of the Compact, Nonthermal Radio Source in the Galactic Center, Sagittarius A*. *Astrophys. J.*, **524**, 805-815 (1999)
 13. Salati, P. & Silk, J. A Stellar Probe of Dark Matter Annihilation in Galactic Nuclei. *Astrophys. J.*, **338**, 24-31 (1989)
 14. Tsiklauri, D. & Viollier, R.D. Dark Matter Concentration in the Galactic Center. *Astrophys. J.*, **500**, 591-595 (1998)
 15. Gondolo, P. & Silk, J. Dark matter annihilation at the Galactic Center. *Phys. Rev. Lett.*, **83**, 1719-1722 (1999)
 16. Torres, D. F., Capozziello, S., & Lambiase, G. A supermassive boson star at the galactic center? *Astrophys. J.*, in press (2000)
 17. Romanowsky, A., and Kochanek, C. to appear in proceedings of “Dynamics of Star Clusters and the Milky Way” (2000)
 18. Sanders, R. H. The case against a massive black hole at the Galactic Centre. *Nature*, **359**, 131-132 (1992)
 19. Morris, M., Ghez, A.M., & Becklin, E.E. The galactic center black hole: clues for the evolution of black holes in galactic nuclei. *Advances in Space Research*, **23**, 959-968 (1999)
 20. Gerhard, O. The Galactic Center He I Stars: Remains of a Dissolved Young Cluster? *Astrophys. J. Lett.*, submitted (2000)

21. Salim, S., & Gould, A. Sagittarius A* “Visual Binaries”: A Direct Measurement of the Galactocentric Distance. *Astrophys. J.*, **523**, 633-641 (1999)
22. Babu, G. J., & Feigelson, E. D. Astrostatistics. *Chapman & Hall*, London, UK (1996).

Acknowledgments This work was supported by the National Science Foundation and the Packard Foundation. We are grateful to James Larkin for swapping telescope time; Ortwin Gerhard, Mike Jura, and Alycia Weinberger for useful input; and telescope observing assistances Joel Aycock, Teresa Chelminiak, Gary Puniwai, Chuck Sorenson, Wayne Wack, Meg Whittle, and software/instrument specialists Al Conrad and Bob Goodrich for their help during the observations. The data presented here were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

Correspondence and requests for materials should be addressed to A.G. (e-mail: ghez@astro.ucla.edu).

Table 1. Measurements for Stars with Significant Accelerations

	S0-1 (S1)	S0-2 (S2)	S0-4 (S8)
Radius from Sgr A*-Magnitude (milli-pc)	4.42 ± 0.03	5.83 ± 0.04	13.15 ± 0.04
Radius from Sgr A*-Position Angle (degrees)	290.1 ± 0.7	3.1 ± 0.4	117.3 ± 0.2
Velocity-Magnitude (km/sec)	1350 ± 40	560 ± 20	990 ± 30
Velocity-Position Angle (degrees)	168 ± 2	241 ± 2	129 ± 2
Acceleration-Magnitude (milli-arcsec/yr ²)	2.4 ± 0.7	5.4 ± 0.3	3.2 ± 0.5
Acceleration-Position Angle (degrees)	80 ± 15	154 ± 4	294 ± 7

Note. — 1 - The primary nomenclature adopted here and in the text is from Ghez et al. (1998); in parentheses star names from Eckart & Genzel (1997) are also given; 2 - all quantities listed are derived from the second order polynomial fit of the data and are given for epoch 1995.53; 3 - all position angles are measured East of North; 4 - all uncertainties are 1σ and are determined by the jackknife resampling method²²; 5 - the radius uncertainties listed include only the relative positional uncertainties and not the uncertainty in the origin used (the nominal position of Sgr A*; see text for offset to measured dynamical center).

Fig. 1.— East-West [*left*] and North-South [*right*] positional offsets from the nominal location of Sgr A* vs. time for S0-1 [*top*], S0-2 [*middle*], & S0-4 [*bottom*]. The offset range shown is scaled to the points in each plot and therefore varies from $\sim 0.''04$ to $\sim 0.''15$. Each of these stars is located with a precision of $\sim 1 - 5$ milli-arcseconds in the individual maps and the alignment of these positions, which is carried out by minimizing the net displacement of all stars as described in Ghez et al.⁴, has an uncertainty of ~ 3 milli-arcseconds based on the half sample bootstrap resampling method²² (the alignment uncertainty is at a minimum at the center of the map and grows linearly with radius). These two uncertainty terms are added in quadrature; the results are used in the fitting process and depicted as errorbars here. In each plot, the solid line shows the second order polynomial used to derive the acceleration term. These plots demonstrate that we are able to measure, for the first time, accelerations of 2 to 5 milli-arcsecond/yr² ($0.3\text{-}0.6 \text{ cm/s}^2$) for stars orbiting a supermassive black hole.

Fig. 2.— The acceleration uncertainty cones and their intersections. The cones' edges represent the directions for which the accelerations deviate by 1σ from their best fit values and their vertices are the time-averaged positions, measured relative to the nominal position of Sgr A*, rather than the positions listed in Table 1. If the accelerations are caused by a single supermassive black hole, or even a spherically symmetric mass distribution, these vectors should intersect at a common location, the center of the mass. The 1σ intersection point lies 0.05 ± 0.03 arcsec East and 0.02 ± 0.03 arcsec South of the nominal position of Sgr A*. The existence of an intersection point suggests that there is indeed a common origin for the acceleration and pinpoints the location of the black hole to within 0.03 arcsec.

Fig. 3.— The minimum enclosed mass implied by each star's acceleration measurement vs. projected distance from the newly determined dynamical center position. If the projection between the plane of the sky and both the radius and the acceleration vectors is θ , then the true mass increases as $1/\cos^3(\theta)$ and the true acceleration and radius vectors increase

as $1/\cos(\theta)$. Also plotted is the mass range implied from a statistical analysis of the ~ 100 velocity vectors that have been measured in the plane of the sky^{3,4}.

Fig. 4.— The measured motion of S0-1, S0-2, & S0-4 and several allowed orbital solutions. Only the measurements obtained in $\sim$ June of each of the 5 years are shown. S0-1 & S0-2 are moving clockwise about Sgr A* and S0-4 is traveling radially outward. In the orbital analysis, two constraints are applied, a central mass of $2.6 \times 10^6 M_\odot$ and a true focus located at the position of Sgr A*. Displayed are orbital solutions with periods of 17, 80, 505 years for S0-2 and 63, 200, 966 years for S0-1.




Figure 1. Ghez et al.




Figure 2. Ghez et al.




Figure 3. Ghez et al.




Figure 4. Ghez et al.